

Package ‘geneNR’

March 10, 2025

Title Automated Gene Identification for Post-GWAS Analysis

Version 1.0.1

Description Facilitates the post-Genome Wide Association Studies (GWAS) analysis of identifying candidate genes within user-defined search window, based on the identified Single Nucleotide Polymorphisms (SNPs) as given by Mazumder AK (2024) <[doi:10.1038/s41598-024-66903-3](https://doi.org/10.1038/s41598-024-66903-3)>. It supports candidate gene analysis for wheat and rice. Just import your GWAS result as explained in the sample_data file and the function does all the manual search and retrieve candidate genes for you, while exporting the results into ready-to-use output.

License CC BY 4.0

Copyright (C) 2025 Nirmalaruban

Encoding UTF-8

RoxygenNote 7.3.2

Depends R (>= 3.5)

LazyData TRUE

Imports readr, stringr, utils, httr, rvest, xml2, writexl

Suggests knitr, rmarkdown, devtools

VignetteBuilder knitr

NeedsCompilation no

Author Rajamani Nirmalaruban [aut, cre, cph],

R. Suvitha [aut],

Rajbir Yadav [aut],

Meda Alekya [aut],

Amit Kumar Mazumder [aut],

Subramani Sugumar [aut],

Prashanth babu [aut],

Manjeet Kumar [aut],

Kiran B Gaikwad [aut],

Naresh Kumar Bainsla [aut],

S. Bhaskar Reddy [aut]

Maintainer Rajamani Nirmalaruban <nirmalaruban97@gmail.com>

Repository CRAN

Date/Publication 2025-03-10 14:50:34 UTC

Contents

geneSNP	2
sample_data_rice	3
sample_data_wheat	3
Index	5

geneSNP	<i>Identifies Candidate Genes based on identified Single Nucleotide Polymorphisms (SNPs) from Genome Wide Association Studies (GWAS) Analysis</i>
---------	---

Description

Identifies Candidate Genes based on identified Single Nucleotide Polymorphisms (SNPs) from Genome Wide Association Studies (GWAS) Analysis

Usage

```
geneSNP(data_file, upstream = 1e+06, downstream = 1e+06, crop = "wheat")
```

Arguments

data_file	The input data in .csv format. (sample_data_wheat or sample_data_rice for demo purpose)
upstream	The search window upstream of the current position of the SNP. (default: 1000000)
downstream	The search window downstream of the current position of the SNP. (default: 1000000)
crop	Either "wheat" or "rice". (default: wheat)

Value

A data frame containing traits, SNP, gene_id, gene_size, and gene_type.

Examples

```
load(system.file("extdata", "precomputed_sample_results.rda", package = "geneNR"))
message(sample_results)

result <- geneSNP("sample_data_wheat", 10000, 10000, crop = "wheat")
result <- geneSNP("sample_data_rice", 10000, 10000, crop = "rice")
```

sample_data_rice	<i>Sample Data</i>
------------------	--------------------

Description

A dataset containing sample data related to genetic markers and associated traits.

Usage

```
sample_data_rice
```

Format

A data frame with columns:

SNP SNP identifier, character.

Chr Chromosome location, character.

Pos Position on the chromosome, numeric.

traits Associated traits, character.

Source

Basha FTM, Sar P, Bhowmick PK, Mahato A, Bisht DS, Iquebal MA, Chakraborty K, Banerjee A, Verma BC, Bhaduri D, Kumar J, Ngangkham U, Saha S, Priyamedha, Mandal NP, Roy S. Genome-wide association study identified QTLs and genes underlying early seedling vigour in aus rice (*Oryza sativa* L.). *Mol Genet Genomics*. 2024 Dec 3;299(1):112. doi: 10.1007/s00438-024-02204-8. PMID: 39625651.

Examples

```
data(sample_data_rice) #lazy loading
```

sample_data_wheat	<i>Sample Data</i>
-------------------	--------------------

Description

A dataset containing sample data related to genetic markers and associated traits.

Usage

```
sample_data_wheat
```

Format

A data frame with columns:

SNP SNP identifier, character.

Chr Chromosome location, character.

Pos Position on the chromosome, numeric.

traits Associated traits, character.

Source

Generated for demonstration purposes

Examples

```
data(sample_data_wheat) #lazy loading
```

Index

* datasets

sample_data_rice, 3

sample_data_wheat, 3

geneSNP, 2

sample_data_rice, 3

sample_data_wheat, 3